

A Prospective Controlled Study of Kidney Donors: Baseline and 6-Month Follow-up

Bertram L. Kasiske, MD,¹ Teresa Anderson-Haag, PharmD, BCPS,¹
 Hassan N. Ibrahim, MD,² Todd E. Pesavento, MD,³ Matthew R. Weir, MD,⁴
 Joseph M. Nogueira, MD,⁴ Fernando G. Cosio, MD,⁵ Edward S. Kraus, MD,⁶
 Hamid H. Rabb, MD,⁶ Roberto S. Kalil, MD,⁷ Andrew A. Posselt, MD,⁸
 Paul L. Kimmel, MD,⁹ and Michael W. Steffes, MD¹⁰

Background: Most previous studies of living kidney donors have been retrospective and have lacked suitable healthy controls. Needed are prospective controlled studies to better understand the effects of a mild reduction in kidney function from kidney donation in otherwise healthy individuals.

Study Design: Prospective, controlled, observational cohort study.

Setting & Participants: Consecutive patients approved for donation at 8 transplant centers in the United States were asked to participate. For every donor enrolled, an equally healthy control with 2 kidneys who theoretically would have been suitable to donate a kidney also was enrolled.

Predictor: Kidney donation.

Measurements: At baseline predonation and at 6 months after donation, medical history, vital signs, measured (iohexol) glomerular filtration rate, and other measurements were collected. There were 201 donors and 198 controls who completed both baseline and 6-month visits and form the basis of this report.

Results: Compared with controls, donors had 28% lower glomerular filtration rates at 6 months (94.6 ± 15.1 [SD] vs 67.6 ± 10.1 mL/min/1.73 m²; $P < 0.001$), associated with 23% greater parathyroid hormone (42.8 ± 15.6 vs 52.7 ± 20.9 pg/mL; $P < 0.001$), 5.4% lower serum phosphate (3.5 ± 0.5 vs 3.3 ± 0.5 mg/dL; $P < 0.001$), 3.7% lower hemoglobin (13.6 ± 1.4 vs 13.1 ± 1.2 g/dL; $P < 0.001$), 8.2% greater uric acid (4.9 ± 1.2 vs 5.3 ± 1.1 mg/dL; $P < 0.001$), 24% greater homocysteine (1.2 ± 0.3 vs 1.5 ± 0.4 mg/L; $P < 0.001$), and 1.5% lower high-density lipoprotein cholesterol (54.9 ± 16.4 vs 54.1 ± 13.9 mg/dL; $P = 0.03$) levels. There were no differences in albumin-creatinine ratios (5.0 [IQR, 4.0-6.6] vs 5.0 [IQR, 3.3-5.4] mg/g; $P = 0.5$), office blood pressures, or glucose homeostasis.

Limitations: Short duration of follow-up and possible bias resulting from an inability to screen controls with kidney and vascular imaging performed in donors.

Conclusions: Kidney donors have some, but not all, abnormalities typically associated with mild chronic kidney disease 6 months after donation. Additional follow-up is warranted.

Am J Kidney Dis. 62(3):577-586. © 2013 by the National Kidney Foundation, Inc.

INDEX WORDS: Chronic kidney disease; unilateral nephrectomy; glomerular filtration rate; patient safety; mineral and bone disorders.

Editorial, p. 448

The shortage of deceased donor kidneys has led to the widespread use of living kidney donors. A number of retrospective studies have reported that short- and long-term outcomes for living kidney donors are excellent.¹ However, these studies have several important limitations. First, they generally fail to

locate all donors who have donated, and the donors who cannot be located may be more likely to have had worse outcomes. Second, many studies do not have a suitable control group. Donors are carefully screened and selected to be healthy, and reports that donors are healthy on follow-up could be biased if donors are compared to the general population or to controls who were not as rigorously screened as donors. The findings that donors live longer than individuals from the

From the ¹Department of Medicine, Hennepin County Medical Center; ²Department of Medicine, University of Minnesota, Minneapolis, MN; ³Department of Medicine, Ohio State University, Columbus, OH; ⁴Department of Medicine, University of Maryland School of Medicine, Baltimore, MD; ⁵Department of Medicine, Mayo Clinic, Rochester, MN; ⁶Department of Medicine, Johns Hopkins University, Baltimore, MD; ⁷Department of Medicine, University of Iowa, Iowa City, IA; ⁸Department of Surgery, University of California at San Francisco, San Francisco, CA; ⁹Division of Kidney Urologic and Hematologic Diseases, National Institute of Diabetes and Digestive and Kidney Diseases, National Insti-

tutes of Health, Bethesda, MD; and ¹⁰Department of Laboratory Medicine and Pathology, University of Minnesota, Minneapolis, MN.

Received October 12, 2012. Accepted in revised form January 18, 2013. Originally published online March 25, 2013.

Address correspondence to Bertram L. Kasiske, MD, Department of Medicine, Hennepin County Medical Center, 701 Park Ave, Minneapolis, MN 55415. E-mail: kasis001@umn.edu

© 2013 by the National Kidney Foundation, Inc.

0272-6386/\$36.00

<http://dx.doi.org/10.1053/j.ajkd.2013.01.027>

general population result from a lack of suitable controls and not from kidney donation prolonging life.^{2,3} Finally, most donors enrolled in retrospective studies donated in an era when selection criteria were more restrictive than they are today.

We designed a multicenter prospective study in which each living donor enrolled with an equally healthy control with 2 kidneys. By including healthy controls, this study permits better assessment of potential harms to kidney donors and thereby provides important information for informing future donors and recipients of the risk of donation. In addition, the present study will allow us to measure parameters in healthy kidney donors that are reported to be abnormal in patients with mild chronic kidney disease (CKD) to determine whether mild reductions in kidney function per se cause abnormalities. In this report, we describe the study and results of baseline and 6-month follow-up visits.

METHODS

Human Subject Protections

Informed consent was obtained from each participant, and the study was approved by the institutional review board at each participating site (University of Minnesota number 0503M67993). In addition, an External Advisory Committee met on August 29, 2006; June 11, 2007; February 20, 2008; December 2, 2009; and May 24, 2011. The External Advisory Committee reviewed the protocol and all revisions and made recommendations regarding the conduct of the study.

Study Design

This prospective observational cohort study was designed and funded for 5 years. Enrollment was to be completed by 2 years, leaving at least 3 years for follow-up. Donors were enrolled after acceptance for donation, but before donation had taken place. For every donor enrolled, one control also was enrolled. Donors and controls completed a baseline predonation visit and visits at 6, 12, 24, and 36 months after donation.

Inclusion Criteria

Initially, individuals selected to be living kidney donors, who were not blood related or who were only distantly blood related to the intended recipient, were asked to participate if they also had a healthy sibling who was willing to participate as a control. The goal was to continue recruitment until 200 pairs of donors and controls had completed baseline and 6-month postdonation visits. The first participant completed a baseline visit July 18, 2006. However, 1 year later, only 23 pairs had enrolled. At that time, the study protocol was amended to enhance enrollment by recruiting any potential living kidney donor. Controls could be any healthy individual who theoretically could be a donor at the study site, not just siblings of enrolled donors. Controls were screened with medical history, vital signs, and basic laboratory tests for kidney disease, but did not undergo other screening tests that may have been performed on donors. Thus, there may have been some bias for donors to be healthier than controls. The last participant completed a baseline study visit February 25, 2011, and enrollment took 4½ years, instead of 2 years as originally planned. Some potential donors completed baseline visits, but did not donate. These donors and their controls were replaced.

Exclusion Criteria

Exclusion criteria for both donors and controls included: (1) unable or unwilling to give informed consent, (2) allergy to intravenous radiocontrast or seafood, and (3) age younger than 18 years. In addition, any living kidney donor exclusion criteria applicable for donors at the study site also were applicable to controls at that site. These generally included evidence of kidney disease (especially proteinuria), invasive cancer, active infection, cardiovascular disease, diabetes, and psychiatric disorders.⁴ Women of child-bearing potential but not pregnant were allowed to participate, but underwent screening for pregnancy before each study visit and were not administered iohexol if/when they became pregnant.

Participating Study Sites

There were initially 7 sites: the University of Minnesota, Minneapolis, MN; Hennepin County Medical Center, Minneapolis, MN; the Mayo Clinic, Rochester, MN; the University of Maryland, Baltimore, MD, the Johns Hopkins School of Medicine, Baltimore, MD, the Ohio State University, Columbus, OH, and the University of Alabama, Birmingham, AL. However, the University of Alabama ended participation before enrolling any participants. The University of Iowa, Iowa City, IA; and the University of California at San Francisco subsequently were added.

Data Collected

Before donation (baseline) and 6 months after donation, participants were evaluated in the Clinical Research Center at each participating site. A complete medical history was obtained. After participants were seated and resting for at least 5 minutes, blood pressure and heart rate were measured 3 times at 1-minute intervals using the right arm (raised to heart level) while they were sitting with feet flat on the floor and resting quietly. Height, body weight, and waist and hip circumference were measured. Blood and urine samples were obtained after an overnight fast. A complete blood count and urinalysis were performed at the site's clinical laboratory. Whole blood, serum, and plasma samples also were sent to the University of Minnesota Advanced Research and Diagnostic Laboratory for fasting glucose, serum electrolytes, calcium, albumin, urea nitrogen, serum creatinine, cystatin C, total cholesterol, high-density lipoprotein cholesterol, calculated low-density lipoprotein cholesterol, triglycerides, lipoprotein(a), homocysteine, glycated hemoglobin A_{1c}, insulin, phosphorus, parathyroid hormone (PTH), uric acid, high-sensitivity C-reactive protein, and fibrinogen analyses and storage at -70°C . A random morning void was used to obtain urine dipstick results. In addition, four 1-mL aliquots of urine were placed into each of 4 clear-capped plastic tubes and frozen: one for urine protein-creatinine and albumin-creatinine ratios and 3 saved specimens, frozen at -70°C . All were transported on dry ice to the Central Laboratory.

Insulin was measured in serum or EDTA plasma on a Roche Elecsys 2010 Analyzer (Roche Diagnostics Corp) using a sandwich immunoassay method (Roche Diagnostics Corp). Intact PTH was measured in serum or EDTA plasma on a Roche Elecsys 2010 Analyzer using a sandwich immunoassay method (Roche Diagnostics). Cystatin C was determined nephelometrically in serum or plasma initially using the Dade Behring BN100 (Dade Behring Inc) nephelometer. In 2012, samples were assayed using the Gentian immunoassay (Gentian AS), which was aligned more closely to an International Federation of Clinical Chemistry and Laboratory Medicine (IFCC) Reference Material (European Reference Material-DA471/IFCC).⁵ All values (including those previously assayed by the Dade-Behring assay) were re-expressed and traceable to the IFCC Reference Material. Total homocysteine was measured in serum or plasma using liquid chromatography (Alli-

Table 1. Enrollment History (number of study pairs)

Study Site	Started ^a	Failed ^b	Enrolled ^c	Dropped ^d	Remained ^e
University of Minnesota	59	-6	53	-1	52
Hennepin County Medical Center	47	-1	46	-2	44
The Ohio State University	34	-2	32	-1	31
University of Maryland	30	-3	27	0	27
The Mayo Clinic, Rochester	23	-3	20	-2	18
Johns Hopkins University	16	-3	13	0	13
University of Iowa	7	0	7	0	7
University of California at San Francisco	6	-1	5	0	5
Total	222	-19	203	-6	197

^aPairs with both completing baseline (predonation) visits.

^bPairs with one or both who did not complete a follow-up visit.

^cPairs with both completing 6-month visits.

^dPairs that dropped out after completing 6-month visits.

^ePairs that remained after completing 6-month visits.

ance 2795 HPLC Separations Module equipped with an autosampler and column oven [Waters Corp]) followed by tandem mass spectrometry.

An iohexol plasma decay method was used to measure glomerular filtration rate (mGFR). A heparin lock was placed in each arm. Slowly, but within 2 minutes, 5 mL of iohexol (Omnipaque; GE Healthcare Inc) was injected, and the catheter was flushed with 10 mL of normal saline solution. Blood samples were drawn from the opposite arm 120, 150, 180, 210, and 240 minutes later, with heparinized plasma separated from each sample. GFR was measured from the plasma decay of iohexol using the Brøchner-Mortensen method.⁶ Iohexol was measured in plasma using a Thermo Scientific SpectraSYSTEM liquid chromatography system (Thermo Separation Products Inc), which consists of a P1000 LC pump coupled with an AS3000 autosampler and UV1000 detector. Chromatographic separation was achieved by means of a Supelcosil LC-18-DB column (Supelco/Sigma-Aldrich Co LLC) with detection at 254 nm. The laboratory interassay coefficient of variation is 2.7%.

CKD Epidemiology Collaboration (CKD-EPI) equations were used to calculate estimated GFR (eGFR), specifically the CKD-EPI creatinine equation from 2009 (yielding eGFR_{cr}),⁷ as well as the CKD-EPI cystatin C and CKD-EPI creatinine-cystatin C equations⁸ published in 2012 (generating eGFR_{cys} and eGFR_{cr-cys}, respectively).

Statistical Analysis

The primary end point of this study is designated to be the difference between donors and controls in the slope of the mGFR measured by iohexol clearance between 6 and 36 months after donations. We estimated that to have 80% power to detect a within-donor difference between donors and controls of 5% in mGFR (~3.25 mL/min/1.73 m²), the study would require 195 donors using a 2-sided paired *t* test and assuming a correlation of 0.6 between measurements from the same individual. This determination was used to select the sample size of the study.

Differences between groups and among visits were assessed using analysis of variance with repeated measures (generalized linear mixed-effects models). This analysis assessed the independent effects of donors versus controls, visits at baseline versus 6 months postdonation, and the interaction between these 2 effects. Results were considered statistically significant for *P* < 0.05, although consideration should be given to the fact that *P* values

were not adjusted for multiple comparisons. Variables that were not normally distributed were logarithmically transformed before analysis. Differences in categorical variables between groups and among visits were assessed with χ^2 . All analyses were carried out with SAS, version 9.2, for the personal computer (SAS Institute Inc).

RESULTS

Enrollment and Study Visits

Two hundred twenty-two pairs started the study and completed the baseline predonation visit (Table 1). Some potential donors did not donate. In 19 pairs, one or both failed to complete a postdonation follow-up visit. However, 203 pairs completed at least one follow-up visit. In 5 cases (2 donors and 3 controls), the 6-month visit was missed and the first follow-up visit was at 12 months. An additional 6 pairs discontinued after completing 6-month follow-up visits. Midway through the study, a decision was made to allow one of a pair to continue in the study if the other decided to discontinue, and to date, 2 controls have dropped out despite their paired donors continuing. Thus, as of June 1, 2012, there were 203 donors and 201 controls still actively participating. However, in this report, we include only baseline and 6-month visits. All anticipated baseline and 6-month visits have been completed (Table 2).

Participant Characteristics at Enrollment

Donor demographics were similar to those of controls (Table 3). However, donors differed from donors in the United States as a whole. In particular, fewer African American and other minority donors enrolled in the study compared with donors in the United States. Ninety-five percent of study donors were white compared to 70% of donors across the United States. Some of this difference can be explained by the

Table 2. Visits Completed as of July 1, 2012

Visit	Controls (n = 201)	Donors (n = 203)
Baseline	201 (100)	203 (100)
6 mo	198 (98.5)	201 (99.0)
12 mo	192 (95.5)	198 (97.5)
24 mo	157 (78.1)	160 (78.8)
36 mo	108 (53.7)	110 (54.2)

Note: Values are given as number (percentage).

location of the study sites. In the states where sites were located, 86% of living kidney donors were white (Table 3). Study participants were more likely to be women and slightly older than living kidney donors in the states where the study sites were located and in the United States. Mean donor age was 43.4 ± 11.9 (SD) years compared to 43.1 ± 11.9 years for controls ($P = 0.8$). Donors and controls had similar medical histories, tobacco use histories, family histories, and medication use at baseline (Table 4). Only 31% of donors had a first-degree relative with CKD, but this was twice that of controls (Table 4).

Blood Pressure, Heart Rate, and Body Size

Heart rates and systolic blood pressures were not different between donors and controls, both declining slightly between the baseline and 6-month visits (Table 5). There were no differences between donors and controls in diastolic blood pressures. There was no statistically significant difference in body weights between donors and controls at 6 months ($P = 0.06$). Waist circumference declined in donors versus controls at 6 months ($P = 0.02$), but there were no differences in body mass index between groups or visits. Approximately 20% of donors and controls

Table 3. Demographic Characteristics of Study Participants Compared to Regional and US Donors During the Same Period (2006-2010)

	Controls (n = 201)	Donors (n = 203)	Study States ^a	United States ^b
Male sex	32.3	32.0	40.3	39.8
White ethnicity	95.0	94.6	85.7	69.7
Age				
18-34 y	30.9	28.1	33.4	33.2
35-49 y	35.3	40.9	42.8	44.6
50-64 y	31.3	28.6	21.9	20.9
≥ 65 y	2.5	2.5	1.8	1.3

Note: Values are given as percentages.

^aRestricted to living kidney donors in the states with participating sites (Minnesota, Ohio, Maryland, Iowa, and California), weighted by the proportion of donors enrolled in those states.

^bAll living donors in the United States who donated, 2006-2010.

were obese at the time of enrollment, with no difference between donors and controls (Table 6).

Kidney Function

mGFR was 28% lower 6 months after donation (Table 6). Neither urine total protein nor urine albumin was affected by donation at 6 months of follow-up.

Laboratory Parameters

Serum albumin concentrations were slightly higher among donors compared with controls at baseline, but declined more in donors than controls after surgery (Table 7). C-Reactive protein and fibrinogen concentrations were not affected by donation. Uric acid and homocysteine concentrations increased in kidney donors. Hemoglobin concentrations were significantly

Table 4. Participant Characteristics at Baseline

Participant Characteristic	Controls (n = 201)	Donors (n = 203)	P ^a
Past medical history			
Hypertension	9 (4.5)	6 (3.0)	0.4
Hyperlipidemia	7 (3.5)	7 (3.5)	0.9
Diabetes mellitus	0 (0.0)	0 (0.0)	0.3
Coronary heart disease	0 (0.0)	1 (0.5)	0.5
Cerebral vascular accident	0 (0.0)	0 (0.0)	0.9
Chronic kidney disease	0 (0.0)	0 (0.0)	0.9
Tobacco use			0.8
Never	132 (65.7)	139 (68.5)	
Former	45 (22.4)	40 (19.7)	
Current	24 (11.9)	24 (11.8)	
Medical history of a parent, sibling, or child			
Chronic kidney disease	30 (14.9)	63 (31.0)	<0.001
Hypertension	90 (44.8)	100 (49.3)	0.4
Diabetes	54 (26.9)	66 (32.5)	0.2
Acute myocardial infarction	40 (19.9)	41 (20.2)	0.9
Coronary revascularization	44 (21.9)	42 (20.7)	0.8
Cerebral vascular accident	27 (13.4)	28 (13.8)	0.9
Cardiovascular disease	77 (38.3)	78 (38.4)	0.9
Medication			
Antidepressant	26 (12.9)	27 (13.3)	0.9
Lipid-lowering agent	31 (15.4)	23 (11.3)	0.2
Antihypertensive agent	10 (5.0)	10 (4.9)	0.9
Aspirin	14 (7.0)	16 (7.9)	0.7
Nonsteroidal anti-inflammatory agent	10 (5.0)	6 (3.0)	0.3
Thyroid replacement	8 (4.0)	11 (5.4)	0.5
Hormone replacement or birth control ^b	29 (21.8)	16 (11.6)	0.02
Vitamin(s)	49 (24.4)	50 (24.6)	0.9
Calcium supplement	18 (9.0)	9 (4.4)	0.07
Other	58 (28.9)	69 (34.0)	0.3

Note: Values are given as number (percentage).

^aBy χ^2 , or (if <10/cell) Fisher exact test.

^bAmong women only.

Table 5. Blood Pressure, Heart Rate, and Body Size

Variable	Baseline Visit		6-mo Visit		<i>P</i> ^a		
	Controls (n = 201)	Donors (n = 203)	Controls (n = 198)	Donors (n = 201)	Controls vs Donors ^b	Baseline vs 6 mo ^c	Inter- action ^d
Heart rate (beats/min)	68.0 ± 9.9 (n = 201)	67.6 ± 10.5 (n = 194)	66.2 ± 10.0 (n = 198)	66.4 ± 10.2 (n = 200)	0.6	0.003	0.5
Systolic blood pressure (mm Hg)	117 ± 13 (n = 201)	117 ± 12 (n = 198)	116 ± 12 (n = 198)	115 ± 11 (n = 199)	0.8	0.003	0.3
Diastolic blood pressure (mm Hg)	70.4 ± 9.0 (n = 201)	70.3 ± 8.8 (n = 198)	70.0 ± 8.5 (n = 198)	70.3 ± 8.5 (n = 199)	0.9	0.6	0.5
Body weight (kg)	77.7 ± 17.1 (n = 199)	77.0 ± 14.8 (n = 199)	78.0 ± 17.3 (n = 197)	76.8 ± 15.2 (n = 199)	0.6	0.6	0.06
Body mass index (kg/m ²)	26.9 ± 5.1 (n = 199)	26.8 ± 4.2 (n = 199)	27.0 ± 5.3 (n = 197)	26.8 ± 4.3 (n = 199)	0.8	0.3	0.3
Waist circumference (cm)	87.3 ± 12.8 (n = 181)	88.0 ± 12.2 (n = 175)	88.0 ± 13.6 (n = 179)	87.2 ± 12.1 (n = 181)	0.7	0.9	0.02

Note: Values are given as mean ± standard deviation (number analyzed). Numbers smaller than 202 reflect missing values.

^aAnalysis of variance with repeated measures. Each variable was analyzed separately and no adjustment was made for multiple comparisons.

^bControls versus donors *P* values test overall differences between donors and controls.

^cBaseline versus 6-month *P* values test overall differences between baseline (predonation) and 6-month visits.

^dInteraction *P* values test the interaction between donors versus controls and baseline versus 6-month visits.

lower in donors at the 6-month visit. Total cholesterol and low-density lipoprotein cholesterol levels were unaffected by donation. Although high-density lipoprotein cholesterol levels were reduced significantly, changes in triglyceride concentrations were not statistically significant. Lipoprotein(a) concentrations did not change with donation. PTH levels increased in donors, whereas serum phosphorus levels declined slightly. Serum total calcium concentrations did not change. Donation had no effect on serum sodium, potassium, chloride, or carbon dioxide levels (data not shown).

DISCUSSION

There have been many studies of kidney donors, but very few have been truly prospective and even fewer have had a suitable control group. Garg et al¹ conducted a systematic review of studies with at least

10 adult kidney donors followed up at least 1 year after donation that measured kidney function and/or urine protein excretion. They identified 48 studies published from 1973 to November 2005. However, only 10 (21%) studies followed donors prospectively; only one of these had a suitable control group (and even this study⁹ did not follow controls prospectively from the time of donation). We searched MEDLINE and EMBASE for studies published since the systematic review of Garg et al.¹ We excluded duplicate publications and located 21 new studies.^{2,10-29} However, only 2 were prospective and neither had a control group.^{13,21}

The controls in the present study were comparable to the kidney donors by all parameters measured (Tables 3-7). Unfortunately, there were few ethnic minorities among donors and controls. Hence, results of this study may not be applicable to other populations. In particular, concerns about the effects of reduced GFR among African American kidney donors cannot be addressed by this study.^{16,30,31} Similarly, few participants in this study were obese.

There are a number of important findings in this study. The increase in PTH levels is in keeping with the correlation between PTH level and kidney function reported in patients with mild CKD. Serum total calcium concentrations were unchanged, whereas serum phosphorus levels were reduced. In 1975, Pabico et al³² noted decreased tubular reabsorption of phosphorus at 10-14 days after nephrectomy in 7 healthy

Table 6. Obesity Among Controls and Donors at Enrollment

Obesity Index ^a	Controls (n = 201)	Donors (n = 203)
Normal, <25.0 kg/m ²	84 (41.8)	72 (35.5)
Overweight, 25.0-29.9 kg/m ²	70 (34.8)	83 (40.9)
Obese, 30.0-39.9 kg/m ²	41 (20.4)	42 (20.7)
Massively obese, ≥40.0 kg/m ²	4 (2.0)	2 (1.0)
Missing/could not be calculated	2 (1.0)	4 (2.0)

Note: Values are given as number (percentage). *P* = 0.5 by χ^2 .

^aBody mass index given after the comma.

Table 7. Laboratory Values

Variable	Baseline Visit		6-mo Visit		P ^a		
	Controls	Donors	Controls	Donors	Controls vs Donors ^b	Baseline vs 6 mo ^c	Interaction ^d
mGFR (mL/min)	106.5 ± 19.3 (n = 186)	106.7 ± 18.6 (n = 181)	104.9 ± 20.2 (n = 194)	74.3 ± 12.9 (n = 193)	0.8	<0.001	<0.001
mGFR (mL/min/1.73 m ²)	96.9 ± 15.3 (n = 186)	96.9 ± 15.3 (n = 181)	94.6 ± 15.1 (n = 194)	67.6 ± 10.1 (n = 193)	0.5	<0.001	<0.001
SCr (mg/dL)	0.79 ± 0.15 (n = 200)	0.80 ± 0.15 (n = 199)	0.80 ± 0.17 (n = 198)	1.16 ± 0.22 (n = 199)	0.8	<0.001	<0.001
eGFR _{cr} (mL/min/1.73 m ²)	100.1 ± 16.0 (n = 200)	99.2 ± 14.4 (n = 199)	99.0 ± 16.0 (n = 198)	65.5 ± 13.1 (n = 199)	0.6	<0.001	<0.001
CysC (mg/dL)	0.81 ± 0.14 (n = 198)	0.80 ± 0.12 (n = 180)	0.81 ± 0.14 (n = 198)	1.11 ± 0.17 (n = 199)	0.6	<0.001	<0.001
eGFR _{cys} (mL/min/1.73 m ²)	102.8 ± 17.6 (n = 198)	103.2 ± 15.4 (n = 180)	102.1 ± 17.5 (n = 198)	71.6 ± 15.3 (n = 199)	0.7	<0.001	<0.001
eGFR _{cr-cys} (mL/min/1.73 m ²)	102.0 ± 16.3 (n = 198)	102.0 ± 13.9 (n = 180)	101.3 ± 16.8 (n = 198)	67.4 ± 11.6 (n = 198)	0.8	<0.001	<0.001
Urea nitrogen (mg/dL)	14.3 ± 3.8 (n = 199)	14.0 ± 3.3 (n = 181)	14.5 ± 4.0 (n = 198)	18.0 ± 4.4 (n = 200)	0.2	<0.001	<0.001
UPCR (g/g)	61 [50-114] (n = 196)	66 [50-128] (n = 175)	62 [50-128] (n = 195)	70 [50-116] (n = 201)	0.3 ^e	0.9 ^e	0.5 ^e
UACR (mg/g)	5.0 [4.0-6.9] (n = 186)	5.0 [3.8-5.8] (n = 167)	5.0 [4.0-6.6] (n = 193)	5.0 [3.3-5.4] (n = 198)	0.07 ^e	0.1 ^e	0.5 ^e
Hemoglobin (g/dL)	13.6 ± 1.2 (n = 194)	13.6 ± 1.2 (n = 198)	13.6 ± 1.4 (n = 193)	13.1 ± 1.2 (n = 194)	0.9	<0.001	<0.001
Leukocyte count (/μL)	6.1 ± 1.6 (n = 195)	5.9 ± 2.0 (n = 198)	6.1 ± 1.7 (n = 193)	5.7 ± 1.5 (n = 194)	0.3	0.2	0.4
Serum albumin (mg/dL)	4.08 ± 0.28 (n = 199)	4.18 ± 0.29 (n = 199)	4.07 ± 0.33 (n = 198)	4.06 ± 0.31 (n = 200)	0.002	<0.001	<0.001
CRP (mg/dL)	1.1 [0.5-2.7] (n = 199)	0.9 [0.4-1.7] (n = 182)	1.4 [0.6-3.1] (n = 198)	1.2 [0.7-2.9] (n = 199)	0.1 ^e	<0.001 ^e	0.2 ^e
Fibrinogen (mg/dL)	295 ± 69 (n = 197)	292 ± 64 (n = 181)	305 ± 67 (n = 198)	300 ± 72 (n = 198)	0.8	0.004	0.7
Homocysteine (mg/L)	1.20 ± 0.35 (n = 193)	1.22 ± 0.39 (n = 176)	1.20 ± 0.34 (n = 196)	1.49 ± 0.43 (n = 198)	0.8	<0.001	<0.001
Uric acid (mg/dL)	4.8 ± 1.1 (n = 200)	4.6 ± 1.1 (n = 198)	4.9 ± 1.2 (n = 198)	5.3 ± 1.1 (n = 200)	0.08	<0.001	<0.001
Serum calcium (mg/dL)	9.16 ± 0.38 (n = 200)	9.26 ± 0.38 (n = 199)	9.19 ± 0.38 (n = 198)	9.24 ± 0.42 (n = 200)	0.02	0.8	0.4
Serum phosphorus (mg/dL)	3.49 ± 0.52 (n = 198)	3.52 ± 0.50 (n = 199)	3.49 ± 0.48 (n = 198)	3.30 ± 0.48 (n = 200)	0.5	<0.001	<0.001
PTH (pg/mL)	42.8 ± 16.3 (n = 199)	42.3 ± 17.8 (n = 180)	42.8 ± 15.6 (n = 198)	52.7 ± 20.9 (n = 200)	0.6	<0.001	<0.001
Cholesterol (mg/dL)	186 ± 37 (n = 200)	185 ± 35 (n = 198)	186 ± 36 (n = 197)	186 ± 35 (n = 199)	0.7	0.7	0.6
LDL cholesterol (mg/dL)	112 ± 33 (n = 198)	110 ± 31 (n = 196)	111 ± 30 (n = 193)	110 ± 31 (n = 193)	0.6	0.7	0.6
HDL cholesterol (mg/dL)	55.2 ± 16.5 (n = 200)	56.2 ± 14.5 (n = 198)	54.9 ± 16.4 (n = 198)	54.1 ± 13.9 (n = 197)	0.5	0.002	0.03
Triglycerides (mg/dL)	77 [55-113] (n = 200)	76 [57-111] (n = 198)	80 [59-119] (n = 197)	84 [64-124] (n = 199)	0.8 ^e	<0.001 ^e	0.05 ^e
Lipoprotein(a) (mg/dL)	12.0 [5-41] (n = 199)	16.0 [5-49] (n = 181)	16.0 [5-43] (n = 198)	20 [5-55] (n = 200)	0.3 ^e	0.003 ^e	0.5 ^e
Hemoglobin A _{1c} (%)	5.3 ± 0.34 (n = 196)	5.2 ± 0.32 (n = 177)	5.3 ± 0.36 (n = 195)	5.3 ± 0.31 (n = 197)	0.03	<0.001	0.6

(Continued)

Table 7 (Cont'd). Laboratory Values

Variable	Baseline Visit		6-mo Visit		<i>P</i> ^a		
	Controls	Donors	Controls	Donors	Controls vs Donors ^b	Baseline vs 6 mo ^c	Interaction ^d
Glucose (mg/dL)	91.8 ± 9.9 (n = 200)	91.9 ± 12.2 (n = 198)	91.2 ± 8.9 (n = 197)	89.2 ± 8.5 (n = 199)	0.9	0.003	0.05
Insulin (pmol/L)	36 [24-60.0] (n = 191)	36 [24-54] (n = 176)	40 [25-64] (n = 192)	36 [24-54] (n = 198)	0.4 ^e	0.01 ^e	0.08 ^e
HOMA IR	1.3 [0.89-2.3] (n = 191)	1.3 [0.8-2.3] (n = 175)	1.5 [1.0-2.5] (n = 191)	1.3 [0.9-2.1] (n = 197)	0.6 ^e	0.04 ^e	0.07 ^e

Note: Values are given as mean ± standard deviation or median [interquartile range] (number sampled). Conversion factors for units: creatinine in mg/dL to μmol/L, ×88.4; urea nitrogen in mg/dL to mmol/L, ×0.357; fibrinogen in mg/dL to μmol/L, ×0.0294; homocysteine in mg/L to μmol/L, ×7.397; uric acid in mg/dL to μmol/L, ×59.48; calcium in mg/dL to mmol/L, ×0.2495; phosphorus in mg/dL to mmol/L, ×0.3229; cholesterol, LDL cholesterol, and HDL cholesterol in mg/dL to mmol/L, ×0.02586; triglycerides in mg/dL to mmol/L, ×0.01129; glucose in mg/dL to mmol/L, ×0.05551.

Abbreviations and definitions: CRP, C-reactive protein; CysC, cystatin C; eGFR_{cr-cys}, glomerular filtration rate estimated by the Chronic Kidney Disease Epidemiology Collaboration creatinine equation; eGFR_{cr-cys}, glomerular filtration rate estimated by Chronic Kidney Disease Epidemiology Collaboration creatinine-cystatin C equation; eGFR_{cys}, glomerular filtration rate estimated by Chronic Kidney Disease Epidemiology Collaboration cystatin C equation; HDL, high-density lipoprotein; HOMA IR, homeostasis model assessment of insulin resistance, where log[HOMA IR] = log[(insulin × glucose)/22.5], with insulin in μU/mL and glucose in mmol/L; LDL, low-density lipoprotein; mGFR, measured glomerular filtration rate (by iohexol plasma clearance); PTH, parathyroid hormone; SCr, serum creatinine; UACR, urine albumin-creatinine ratio; UPCR, urine protein-creatinine ratio.

^aAnalysis of variance with repeated measures. Each variable was analyzed separately and no adjustment was made for multiple comparisons. Values not normally distributed were logarithmically transformed before analysis.

^bControls versus donors *P* values test overall differences between donors and controls.

^cBaseline versus 6-month *P* values test overall differences between baseline (predonation) and 6-month visits.

^dInteraction *P* values test the interaction between donors versus controls and baseline versus 6-month visits.

^eBased on logarithmically transformed values.

donors. Friedlander et al³³ found that 17 donors had reduced tubular reabsorption of phosphate at 1 and 6 months after donation, and this was associated with an increase in carboxyl-terminal PTH levels. Other cross-sectional studies have reported reduced 1,25-dihydroxyvitamin D³⁴ and increased fibroblast growth factor-23 (FGF-23) levels.³⁵ In a recent cross-sectional study, Young et al³⁶ found increased FGF-23 levels, which correlated with the decline in eGFRs in donors. In the Young et al³⁶ study, vitamin D and serum phosphate concentrations were reduced, whereas fractional excretion of phosphate was increased compared with healthy controls. In the present study, vitamin D and FGF-23 were not measured. Additional longitudinal studies are needed to determine how the decline in kidney function after donation leads to elevated PTH and reduced phosphorus concentrations, and what role vitamin D and FGF-23 play in these alterations.

It previously has been reported that patients with CKD have abnormalities in lipoprotein metabolism, including elevated triglyceride and reduced high-density lipoprotein cholesterol levels.^{37,38} However, the extent to which these lipid alterations were due to reduced kidney function per se or to the underlying causes of CKD has been unclear from uncontrolled observational studies. The present study indicates that the mild reduction in kidney function resulting from donor nephrectomy caused a decline in high-density lipoprotein cholesterol levels, but no statistically sig-

nificant changes in triglyceride, low-density lipoprotein cholesterol, or lipoprotein(a) levels.

Hyperuricemia has long been suggested to cause CKD,³⁹⁻⁴³ hypertension,⁴⁴ diabetes,⁴⁵ and cardiovascular disease.⁴⁶⁻⁴⁸ However, this has been a source of ongoing controversy because it is possible that hypertension, diabetes, and cardiovascular disease directly or indirectly cause hyperuricemia, resulting in so-called reverse causation. The present study unequivocally shows that a reduction in GFR causes a significant increase in serum uric acid level, even in otherwise healthy individuals. Thus, whatever other factors may be causing an association between CKD and hyperuricemia, the reduction in GFR itself may explain much or all of the observed association.

Cross-sectional studies suggest that CKD is associated with abnormalities in glucose homeostasis and insulin resistance.⁴⁹⁻⁵⁴ However, it is again unclear whether altered kidney function or underlying causes of CKD result in these abnormalities. In the present study, there were no differences between donors and controls in fasting glucose, hemoglobin A_{1c}, or insulin concentrations or the calculated homeostasis model assessment of insulin resistance. These data suggest that short-term mild reductions in kidney function resulting from kidney donation do not cause alterations in glucose homeostasis or insulin resistance.

In the present study, hemoglobin levels were significantly lower in donors 6 months after donation compared with controls (Table 7). We are not aware of this

previously being reported. Possible explanations include mild anemia due to surgical blood loss, iron deficiency, and/or reduced erythropoietin due to the reduced kidney function. Whether this resolves with time remains to be determined.

Cross-sectional observational studies have shown a correlation between reduced kidney function and increased homocysteine concentrations.⁵⁵⁻⁵⁷ The present study provides conclusive evidence that a mild reduction in kidney function causes increases in homocysteine concentrations. Nevertheless, recent randomized trials have failed to show that reducing homocysteine levels improves patient outcomes.^{58,59} It therefore is unclear whether an elevated homocysteine level is harmful.

Several studies have suggested that kidney donors have mild proteinuria.¹ It therefore is interesting that there were no differences between donors and controls in urine total protein or albumin-creatinine ratios in the present study. Similarly, a number of investigations have suggested that blood pressure is mildly elevated in kidney donors.⁶⁰ However, blood pressure in the present study was not different between donors versus controls. The lack of differences in protein excretion and blood pressure could reflect the short-term follow-up in the present study.

Although there were statistically significant effects of donation on serum albumin concentrations, these effects were small and difficult to interpret (Table 7). At baseline, serum albumin concentrations were numerically higher in donors than controls, declining to control levels at 6 months. It is difficult to conclude with certainty that donation caused a significant reduction in serum albumin concentrations. Additional follow-up may help clarify these changes. It is noteworthy that several other biomarkers of inflammation were not affected by donation. In particular, C-reactive protein and fibrinogen concentrations were unchanged after donation (Table 7). These results, if sustained over time, may suggest that the putative inflammatory state associated with mild CKD may not be the result of a decline in kidney function as much as the underlying disease and the causes of CKD.

The decline in mGFR at 6 months was similar to that reported by others using various inert markers of GFR.⁶¹⁻⁷⁰ It is noteworthy that estimating equations using creatinine and/or cystatin C produced results similar to those of the pre- and postdonation mGFRs. These equations slightly overestimated mGFR before donation and mGFR in controls with normal kidney function, but this bias was less apparent among donors after donation (Table 7). Long-term follow-up may help determine whether changes in eGFR accu-

rately reflect changes in GFR measured by iohexol, and which formulation may be most accurate.

There are some important limitations to this study. The controls could not be subjected to the same rigorous screening process that donors underwent, and therefore it is possible that controls were healthier than donors in ways not reflected by the study measurements reported here. However, more donors (31%) than controls (15%) were blood relatives of individuals with CKD (Table 4), which could predispose the donors to CKD. Finally, a relatively large number of variables were examined, making it possible that some differences in results were due to chance (type I statistical error). Therefore, it will be important to confirm these results in other studies, if possible.

In summary, the short-term results of this study demonstrate that a number of physiologic changes associated with CKD are found in donors with mild declines in GFR. However, a number of the reported changes wrought by CKD, such as increased blood pressure, were not found in kidney donors. This ongoing study offers a unique opportunity to examine the effects of donation with its mild to moderate reductions in GFR on parameters of interest in patients with CKD. It thereby will help us understand the consequences of CKD, as well as the safety of kidney donation.

ACKNOWLEDGEMENTS

The authors thank Jon Snyder, PhD, for advice in performing the statistical analysis.

Support: This study was funded by the National Institutes of Health under cooperative agreement U01 DK066013, Paul L. Kimmel, MD, Project Officer. It was also supported by funds from the Minneapolis Medical Research Foundation, Minneapolis, MN.

Financial Disclosure: The authors declare that they have no other relevant financial interests.

REFERENCES

1. Garg AX, Muirhead N, Knoll G, et al. Proteinuria and reduced kidney function in living kidney donors: a systematic review, meta-analysis, and meta-regression. *Kidney Int.* 2006; 70(10):1801-1810.
2. Segev DL, Muzaale AD, Caffo BS, et al. Perioperative mortality and long-term survival following live kidney donation. *JAMA.* 2010;303(10):959-966.
3. Andreoni KA. "Live kidney donors live longer" and would you like to buy part of a bridge in Brooklyn? *Arch Surg.* 2010; 145(7):701-702.
4. The consensus statement of the Amsterdam Forum on the Care of the Live Kidney Donor. *Transplantation.* 2004;78(4):491-492.
5. Inker LA, Eckfeldt J, Levey AS, et al. Expressing the CKD-EPI (Chronic Kidney Disease Epidemiology Collaboration) cystatin C equations for estimating GFR with standardized serum cystatin C values. *Am J Kidney Dis.* 2011;58(4):682-684.
6. Brochner-Mortensen J. A simple method for the determination of glomerular filtration rate. *Scand J Clin Lab Invest.* 1972; 30(3):271-274.

7. Levey AS, Stevens LA, Schmid CH, et al. A new equation to estimate glomerular filtration rate. *Ann Intern Med.* 2009;150(9):604-612.
8. Inker LA, Schmid CH, Tighiouart H, et al. Estimating glomerular filtration rate from serum creatinine and cystatin C. *N Engl J Med.* 2012;367(1):20-29.
9. Dunn JF, Richie RE, MacDonell RC Jr, Nylander WA Jr, Johnson HK, Sawyers JL. Living related kidney donors: a 14-year experience. *Ann Surg.* 1986;203(6):637-643.
10. Gossmann J, Wilhelm A, Kachel HG, et al. Long-term consequences of live kidney donation follow-up in 93% of living kidney donors in a single transplant center. *Am J Transplant.* 2005;5(10):2417-2424.
11. Fehrman-Ekholm I, Norden G, Lennerling A, et al. Incidence of end-stage renal disease among live kidney donors. *Transplantation.* 2006;82(12):1646-1648.
12. Gres P, Avances C, Iborra F, Mourad G, Guiter J. [Long-term morbidity of living donor kidney harvesting]. *Prog Urol.* 2007;17(2):194-198.
13. Minnee RC, Bemelman WA, Polle SW, et al. Older living kidney donors: surgical outcome and quality of life. *Transplantation.* 2008;86(2):251-256.
14. Ibrahim HN, Foley R, Tan L, et al. Long-term consequences of kidney donation. *N Engl J Med.* 2009;360(5):459-469.
15. Albertsmeyer S, Renner FC, Yildiz S, et al. One hundred six live kidney donors in a single German transplantation center: renal, physical, and psychological follow-up. *Transplant Proc.* 2010;42(10):3992-3993.
16. Lentine KL, Schnitzler MA, Xiao H, et al. Racial variation in medical outcomes among living kidney donors. *N Engl J Med.* 2010;363(8):724-732.
17. Okamoto M, Suzuki T, Fujiki M, et al. The consequences for live kidney donors with preexisting glucose intolerance without diabetic complication: analysis at a single Japanese center. *Transplantation.* 2010;89(11):1391-1395.
18. Storsley LJ, Young A, Rush DN, et al. Long-term medical outcomes among Aboriginal living kidney donors. *Transplantation.* 2010;90(4):401-406.
19. Berger JC, Muzaale AD, James N, et al. Living kidney donors ages 70 and older: recipient and donor outcomes. *Clin J Am Soc Nephrol.* 2011;6(12):2887-2893.
20. Cheriakh WS, Young CJ, Kramer BF, Taranto SE, Randall HB, Fan PY. Ethnic and gender related differences in the risk of end-stage renal disease after living kidney donation. *Am J Transplant.* 2011;11(8):1650-1655.
21. Dols LF, Kok NF, Roodnat JJ, et al. Living kidney donors: impact of age on long-term safety. *Am J Transplant.* 2011;11(4):737-742.
22. Wafa EW, Refaie AF, Abbas TM, et al. End-stage renal disease among living-kidney donors: single-center experience. *Exp Clin Transplant.* 2011;9(1):14-19.
23. Chu KH, Poon CK, Lam CM, et al. Long-term outcomes of living kidney donors: a single centre experience of 29 years. *Nephrology (Carlton).* 2012;17(1):85-88.
24. Fournier C, Pallet N, Cherqaoui Z, et al. Very long-term follow-up of living kidney donors. *Transpl Int.* 2012;25(4):385-390.
25. Garg AX, Meirambayeva A, Huang A, et al. Cardiovascular disease in kidney donors: matched cohort study. *BMJ.* 2012;344:e1203.
26. Mjoen G, Reisaeter A, Hallan S, et al. Overall and cardiovascular mortality in Norwegian kidney donors compared to the background population. *Nephrol Dial Transplant.* 2012;27(1):443-447.
27. O'Brien B, Mastoridis S, Sabharwal A, Hakim N, Taube D, Papalois V. Expanding the donor pool: living donor nephrectomy in the elderly and the overweight. *Transplantation.* 2012;93(11):1158-1165.
28. Tent H, Sanders JS, Rook M, et al. Effects of preexistent hypertension on blood pressure and residual renal function after donor nephrectomy. *Transplantation.* 2012;93(4):412-417.
29. Tsai SF, Shu KH, Ho HC, et al. Long-term outcomes of living kidney donors over the past 28 years in a single center in Taiwan. *Transplant Proc.* 2012;44(1):39-42.
30. Gibney EM, King AL, Maluf DG, Garg AX, Parikh CR. Living kidney donors requiring transplantation: focus on African Americans. *Transplantation.* 2007;84(5):647-649.
31. Cohen DM, Mittalhenkle A, Scott DL, Young CJ, Norman DJ. African American living-kidney donors should be screened for APOL1 risk alleles. *Transplantation.* 2011;92(7):722-725.
32. Pabico RC, McKenna BA, Freeman RB. Renal function before and after unilateral nephrectomy in renal donors. *Kidney Int.* 1975;8(3):166-175.
33. Friedlander MA, Lemke JH, Horst RL. The effect of uninephrectomy on mineral metabolism in normal human kidney donors. *Am J Kidney Dis.* 1988;11(5):393-401.
34. Bieniasz M, Kwiatkowski A, Domagala P, et al. Serum concentration of vitamin D and parathyroid hormone after living kidney donation. *Transplant Proc.* 2009;41(8):3067-3068.
35. Westerberg PA, Ljunggren O, Larsson TE, Wadstrom J, Linde T. Fibroblast growth factor-23 and mineral metabolism after unilateral nephrectomy. *Nephrol Dial Transplant.* 2010;25(12):4068-4071.
36. Young A, Hodsman AB, Boudville N, et al. Bone and mineral metabolism and fibroblast growth factor 23 levels after kidney donation. *Am J Kidney Dis.* 2012;59(6):761-769.
37. Kwan BC, Kronenberg F, Beddhu S, Cheung AK. Lipoprotein metabolism and lipid management in chronic kidney disease. *J Am Soc Nephrol.* 2007;18(4):1246-1261.
38. Vaziri ND, Navab M, Fogelman AM. HDL metabolism and activity in chronic kidney disease. *Nat Rev Nephrol.* 2010;6(5):287-296.
39. Iseki K, Oshiro S, Tozawa M, Iseki C, Ikemiya Y, Takishita S. Significance of hyperuricemia in the early detection of renal failure in a cohort of screened subjects. *Hypertens Res.* 2001;24(6):691-697.
40. Obermayr RP, Temml C, Knechtelsdorfer M, et al. Predictors of new-onset decline in kidney function in a general middle-European population. *Nephrol Dial Transplant.* 2008;23(4):1265-1273.
41. Weiner DE, Tighiouart H, Elsayed EF, Griffith JL, Salem DN, Levey AS. Uric acid and incident kidney disease in the community. *J Am Soc Nephrol.* 2008;19(6):1204-1211.
42. Hsu CY, Iribarren C, McCulloch CE, Darbinian J, Go AS. Risk factors for end-stage renal disease: 25-year follow-up. *Arch Intern Med.* 2009;169(4):342-350.
43. Ben-Dov IZ, Kark JD. Serum uric acid is a GFR-independent long-term predictor of acute and chronic renal insufficiency: the Jerusalem Lipid Research Clinic cohort study. *Nephrol Dial Transplant.* 2011;26(8):2558-2566.
44. Grayson PC, Kim SY, LaValley M, Choi HK. Hyperuricemia and incident hypertension: a systematic review and meta-analysis. *Arthritis Care Res (Hoboken).* 2011;63(1):102-110.
45. Kodama S, Saito K, Yachi Y, et al. Association between serum uric acid and development of type 2 diabetes. *Diabetes Care.* 2009;32(9):1737-1742.
46. Kim SY, Guevara JP, Kim KM, Choi HK, Heitjan DF, Albert DA. Hyperuricemia and risk of stroke: a systematic review and meta-analysis. *Arthritis Rheum.* 2009;61(7):885-892.
47. Kim SY, Guevara JP, Kim KM, Choi HK, Heitjan DF, Albert DA. Hyperuricemia and coronary heart disease: a system-

atic review and meta-analysis. *Arthritis Care Res (Hoboken)*. 2010;62(2):170-180.

48. Feig DI, Kang DH, Johnson RJ. Uric acid and cardiovascular risk. *N Engl J Med*. 2008;359(17):1811-1821.

49. Dzurik R, Spustova V, Janekova K. The prevalence of insulin resistance in kidney disease patients before the development of renal failure. *Nephron*. 1995;69(3):281-285.

50. Eidemak I, Feldt-Rasmussen B, Kanstrup IL, Nielsen SL, Schmitz O, Strandgaard S. Insulin resistance and hyperinsulinemia in mild to moderate progressive chronic renal failure and its association with aerobic work capacity. *Diabetologia*. 1995;38(5):565-572.

51. Vareesangthip K, Tong P, Wilkinson R, Thomas TH. Insulin resistance in adult polycystic kidney disease. *Kidney Int*. 1997;52(2):503-508.

52. Fliser D, Pacini G, Engelleiter R, et al. Insulin resistance and hyperinsulinemia are already present in patients with incipient renal disease. *Kidney Int*. 1998;53(5):1343-1347.

53. Kubo M, Kiyohara Y, Kato I, et al. Effect of hyperinsulinemia on renal function in a general Japanese population: the Hisayama Study. *Kidney Int*. 1999;55(6):2450-2456.

54. Kato Y, Hayashi M, Ohno Y, Suzawa T, Sasaki T, Saruta T. Mild renal dysfunction is associated with insulin resistance in chronic glomerulonephritis. *Clin Nephrol*. 2000;54(5):366-373.

55. Norlund L, Grubb A, Fex G, et al. The increase of plasma homocysteine concentrations with age is partly due to the deterioration of renal function as determined by plasma cystatin C. *Clin Chem Lab Med*. 1998;36(3):175-178.

56. Wollesen F, Brattstrom L, Refsum H, Ueland PM, Berglund L, Berne C. Plasma total homocysteine and cysteine in relation to glomerular filtration rate in diabetes mellitus. *Kidney Int*. 1999;55(3):1028-1035.

57. Francis ME, Eggers PW, Hostetter TH, Briggs JP. Association between serum homocysteine and markers of impaired kidney function in adults in the United States. *Kidney Int*. 2004;66(1):303-312.

58. Clarke R, Halsey J, Lewington S, et al. Effects of lowering homocysteine levels with B vitamins on cardiovascular disease, cancer, and cause-specific mortality: meta-analysis of 8 randomized trials involving 37 485 individuals. *Arch Intern Med*. 2010;170(18):1622-1631.

59. Jardine MJ, Kang A, Zoungas S, et al. The effect of folic acid based homocysteine lowering on cardiovascular events in people with kidney disease: systematic review and meta-analysis. *BMJ*. 2012;344:e3533.

60. Boudville N, Prasad GV, Knoll G, et al. Meta-analysis: risk for hypertension in living kidney donors. *Ann Intern Med*. 2006;145(3):185-196.

61. Aurell M, Ewald J. The use of living donors. Glomerular filtration rate during the first year after donor nephrectomy. *Scand J Urol Nephrol Suppl*. 1981;64:137-142.

62. Bertolatus JA, Friedlander MA, Scheidt C, Hunsicker LG. Urinary albumin excretion after donor nephrectomy. *Am J Kidney Dis*. 1985;5(3):165-169.

63. Chan MK. Protein loading test before and after kidney donation. *Aust N Z J Med*. 1986;16(5):691-694.

64. ter Wee PM, Tegzess AM, Donker AJ. Renal reserve filtration capacity before and after kidney donation. *J Intern Med*. 1990;228(4):393-399.

65. Bock HA, Bachofen M, Landmann J, Thiel G. Glomerular hyperfiltration after unilateral nephrectomy in living kidney donors. *Transpl Int*. 1992;5(suppl 1):S156-S159.

66. ter Wee PM, Tegzess AM, Donker AJM. Pair-tested renal reserve filtration capacity in kidney recipients and their donors. *J Am Soc Nephrol*. 1994;4:1798-1808.

67. Shokeir AA, El-Diasty TA, Gad HM, et al. Radioisotopic evaluation of renal function in cyclosporine-treated pediatric and adult renal transplant recipients and their living donors. *Transplantation*. 1994;58:1171-1175.

68. Kamper AL, Holstein-Rathlou NH, Strandgaard S, Leyssac PP, Munck O. Glomerular filtration rate and segmental tubular function in the early phase after transplantation/uninephrectomy in recipients and their living-related kidney donors. *Clin Sci (Lond)*. 1994;87(5):519-523.

69. Velosa JA, Offord KP, Schroeder DR. Effect of age, sex, and glomerular filtration rate on renal function outcome of living kidney donors. *Transplantation*. 1995;60(12):1618-1621.

70. Velosa JA, Griffin MD, Larson TS, et al. Can a transplanted living donor kidney function equivalently to its native partner? *Am J Transplant*. 2002;2(252):259.